No.



9800001

## The Curators of the Unibersity of Missouri

MCCCUS, THERE HAS BEEN PRESENTED TO THE

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT, VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN DUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT  ${
m VARIETY}$ ECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'Maverick'

In Testimony Mexicos, I have hereunto set my hand and caused the seal of the Hant Hariety Protection Office to be affixed at the City of Washington, D.C. this thirty-first day of March, in the year of our Lord two thousand.

REPRODUCE LOCALLY. Include form number and date on all	reproductions.		FORM APPROVED - OMB NO. 0581-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTEC	TION OFFICE	The following statements are ma 1974 (5 U.S.C. 552a) and the Paj	de in accordance with the Privacy Act of perwork Reduction Act (PRA) of 1995.
APPLICATION FOR PLANT VARIETY PROTECTION  (Instructions and information collection burden statement)	CERTIFICATE		to determine if a plant variety protection C. 2421). Information is held confidential 2426).
NAME OF APPLICANT(S) (as it is to appear on the Certificate)	TO T	2. TEMPORARY DESIGNATION OR	3. VARIETY NAME
The Curators of the University of Misso	uri	LN90-4455	Maverick
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Countr University of Missouri	y)	6. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY
311 Jesse Hall Columbia, MO 65211		573-882-3211	9800001
oorama vaş ilo oozii		6. FAX (include area code)	F DATE
		573-884=5446	01/OR+197
	8. FAMILY NAME (Botani	cal)	FILING AND EXAMINATION FEE:
Glycine max (L.) Merr.	Leguminosa	e	* 2450.00
9. CROP KIND NAME (Common name)		National Control of the Control of t	S DATE SILVARION
Soybean			R CI/OCI/9/
10. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATI Educational Organization	ON (corporation, partnersh	ip, association, etc.) (Common name)	C CERTIFICATION FEE:
11. IF INCORPORATED, GIVE STATE OF INCORPORATION		12. DATE OF INCORPORATION	E DATE
Missouri		,	Nov. 8, 1779
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERVE D. A. Sleper	IN THIS APPLICATION A	ND RECEIVE ALL PAPERS	14. TELEPHONE (include area code) 573-882-7320
Department of Agronomy 201 Waters Hall			373-662-7320
University of Missouri			15. FAX (include area code)
Columbia, MO 65211			573-882=1467
16. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow Inst. a.   ■ Exhibit A. Origin and Breeding History of the Variety	ructions on reverse)		
b. 🔀 Exhibit B. Statement of Distinctness			in in the second of the secon
c. X Exhibit C. Objective Description of the Variety	en de la Companya de La Companya de la Com		
d. X Exhibit D. Additional Description of the Variety (Optional)		•	and the second second
e. X Exhibit E. Statement of the Basis of the Applicant's Ownership			The state of the s
f. X Voucher Sample (2,600 viable untreated seeds or, for tuber propagated vg. X Filing and Examination Fee (\$2,450), made payable to "Treasurer of the			ed in an approved public repository)
17. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VA		CLASS OF CERTIFIED SEED? (See Section	n 83(a) of the Plant Variety Protection Acti
18. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS GENERATIONS?	TO NUMBER OF 19.	IF "YES" TO ITEM 18, WHICH CLASSES	OF PRODUCTION BEYOND BREEDER SEED?
☑ YES ☐ NO	and the second	☑ FOUNDATION ☐ REGISTER	
20. HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEN RELEA  YES (If "yes," give names of countries and dates)		R SALE, OR MARKETED IN THE U.S. OR C	THER COUNTRIES?
the state of the s			<u> </u>
21. The applicant(s) declare that a viable sample of basic seed of the variety will be fu applicable, or for a tuber propagated variety a tissue culture will be deposited in a	rnished with application an public repository and main	d will be replenished upon request in accordanced for the duration of the certificate.	dance with such regulations as may be
The undersigned applicant(s) is(are) the owner(s) of this sexually reproduced or tuk Section 42, and is entitled to protection under the provisions of Section 42 of the f	per propagated plant variety	/, and believe(s) that the variety is new, di	stinct, uniform, and stable as required in
Applicant(s) is(are) informed that false representation herein can jeopardize protect	ion and result in penalties.		
SIGNATURE OF APPLICANT (OWNERS)	SIGNATUR	E OF APPLICANT (Owner(s))	443 - 44 November 1997
NAME (Nease print or type)	NAME (Plea	ise print or type	
Roger L. Mitchell			
Director of Ag. Experiment Stat.	CAPACITY	OR TITLE	DATE
STD-470 (03.96) (Provious editions are to be destroyed)	1		f

#### Exhibit A

#### Origin and Breeding History of the Variety

Maverick originated as a F<sub>4</sub>-plant selection from the cross of LN86-4668 x 'Resnik' made at the Illinois Agricultural Experiment Station. LN86-4668 is a selection from the cross 'Fayette' x 'Hardin'. The original cross was made in the field in the summer of 1987, and the  $F_1$  generation grown in the field in 1988. The F<sub>2</sub> and F<sub>3</sub> generations were advanced by single-seed-descent in Puerto Rico during the winter of 1988 andd 1989 and the F<sub>4</sub> generation was grown at Urbana in the summer of 1989. In the greenhouse at the University of Illinois during the winter of 1989 and 1990, progeny from single plants selected in the summer of 1989 were evaluated for resistance to races 3 and 14 of soybean cyst nematode (caused by Heterodera glycines Inchinohe) and resistance to races 1 and 3 of Phytophthora rot (caused by Phytophthora sojae M.J. Kaufmann & J.W. Gerdemann). During the winter of 1995 in the greenhouse at the University of Missouri, single plants of Maverick were evaluated for resistance to races 3 and 14 of the soybean cyst nematode. The F<sub>5</sub> generation was grown as plant rows in 1990 at the University of Illinois. Singleplant rows were selected, composited, and evaluated in replicated yield trials in Illinois in 1991. Maverick was identified and evaluated in replicated tests at multiple-locations in Missouri and Illinois, 1992 through 1995. Maverick was tested in the Preliminary SCN III Test in 1994 and Uniform SCN III Tests in 1995 and 1996 of the Northern Regional Soybean Cyst Nematode Test. Also, it was evaluated in Uniform III Tests in 1996 of the Uniform Soybean Tests-Northern Region Test.

Maverick was stable and uniform from the  $F_5$  generation through our seed increase and testing program through  $F_{11}$  generation. The only variants observed on average were up to 1% variant in flower color and up to 1% variant in hila color.

#### Exhibit B

#### Statement of Distinctness

Maverick most closely resembles Iroquois in plant habit (indeterminate), maturity (Group III), flower color (purple), pubescence color (grey), hypocotyl color (light purple), pod color at maturity (brown), seed size (14 g/100 seeds), and resistance to Phytophthora rot (race 1).

Maverick has dull yellow seeds, buff hila, and resistance to soybean cyst nematode (races 3 and 14) while Iroquois has yellow seeds, imperfect black hila, and is susceptible to the soybean cyst nematode.

#### AGRICULTURAL MARKETING SERVICE

PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MARYLAND 20705

## OBJECTIVE DESCRIPTION OF VARIETY SOYBEAN (Glycine max L.)

				EAN (Glycine			
NAME O	F APPLICANT	(S)	oweitu	TEMPORARY	DESIGNATION VA	RIETY NAME	
of M	issouri	of the Univ	•	LN90-4	455	Maverick	
ADDRES Univ	s <i>(Street and I</i> ersity of	Vo., or R.F.D. No., Missouri	City, State, and Zip Co	ode)		FOR OFF	CIAL USE ONLY
	Universit				P\	PO NUMBER	
	mbia, MO	<del></del>	·			98:	60001
Choose t	the appropriat	te response which	characterizes the v	ariety in the fea	tures described belo	ow. When the nu	mber of significant o
Starred c	haracters + 2	r than the numb	er of boxes provided	1, place a zero in	the first box wher	number is 9 or 1	ess (e.g., 0 9
when infe	ormation is av	vailable.	idamental (O an ade	quate soyoean y	ariety description.	Other characters	should be described
1. SEED S	SHAPE:		$\Theta$ 0				
1			$\Theta$	, 1		t	
لـــٰــا			L W	т	•		
1	1 = Spherical (L	.W, L/T, and T/W	ratios = < 1.2)	2 = Spl	nerical Flattened (L/M	/ ratio > 1.2; L/T r	atio = <b>&lt;</b> 1.2)
	3 - Ciongate (L)	/T ratio > 1.2; T/\	W = ₹ 1.2)	4 ≃ Elo	ngate Flattened (L/T	ratio > 1.2; T/W	> 1.2)
2. SEED C	OAT COLOR:	(Mature Seed)		<u></u>	-		
1 1	= Yellow	2 = Green	3 = Brown	4 ≃ Black			
			<i>y</i> .	4 - Black	5 = Other (Spec	city)	
3. SEED C	OAT LUSTER:	: (Mature Hand Sh	elled Seed)		·····		
1 1	= Dull ('Corso)	y 79'; 'Braxton')	2 = Shiny ('Nebs	oy'; 'Gasoy 17')		• •	12
<del></del>	<del></del>				<u> </u>	Ar La	
7 4. SEED SI	ZE: (Mature S	leed)	3A	·	**		
1 4 G	rams per 100 se	eeds			, P. a		
5. HILUM (	COLOR: (Matu	ıre Seed)			<u>,,,</u>		* :
1 1	= Buff	2 = Yellow	3 = Brown	4 = Gray 5	= Imperfect Black	6 = Black	7 = Other (Specify)
6. COTYLE	DON COLOR:	(Mature Seed) X			· · · · · · · · · · · · · · · · · · ·		
[1]		•					*
	= Yellow 2	2 = Green			• •		
7. SEED PR	OTEIN PEROV	(IDASE ACTIVITY	<del></del>			· · · · · · · · · · · · · · · · · · ·	····
	O'EMIT EROX	NDASE ACTIVITY	r: 🗸	•		· 	
1 =	Low 2	2 = High		1			
9 CEED DO					<del></del>		
o. SCEUPRO	OTEIN ELECT	ROPHORETIC BA	ND: 🗸				
1 =	Type A (SP1 <sup>a</sup> )	)	2 = Type B (SP1 <sup>b</sup> )				
9. НҮРОСОТ	TYL COLOR:	· · · · · · · · · · · · · · · · · · ·			······································		
3 1=	Green only ("E Light Purple be	eiow cotyledons ('E	Beeson'; 'Pickett 71')		w cotyledons (Wood	worth"; "Tracy")	
3 ≥	Dark Purpie ex	tending to unifolia	te leaves ('Hodgson'; '	Coker Hampton 20	66A')		•
3 ≥		tending to unifolia	ite leaves ('Hodgson'; '	Coker Hampton 2	66A*) 	<del></del>	·

FORM LMGS-470-57 (6-83)

11. LEAFLET	SIZE:	3600000
. 5] ¹	= Small ('Amsoy 71'; 'A5312') 2 = Medium ('Corsoy 79'; 'Gasoy 17') = Large ('Crawford'; 'Tracy')	,
12. LEAF COL	OR:	
2 1:	= Light Green ('Weber'; 'York') 2 = Medium Green ('Corsoy 79'; 'Braxton') = Dark Green ('Gnome'; 'Tracy')	<u></u> -
★ 13. FLOWER C	TOLOR:	
2 1-	= White 2 = Purple 3 = White with purple throat	
★ 14. POD COLO	R:	
<u> </u>	Tan 2 = Brown 3 = Błack	
★ 15. PLANT PUB	BESCENCE COLOR:	
<u></u>	Gray 2 = Brown (Tawny)	
16. PLANT TYP	PES:	
1 = 3 =	Slender ('Essex'; 'Amsoy 71')  2 = Intermediate ('Amcor'; 'Braxton')  Bushy ('Gnome'; 'Govan')	<del>.</del>
★ 17. PLANT HAB	BIT:	
3 1 = 3 =	Determinate ('Gnome'; 'Braxton') 2 = Semi-Determinate ('Will') Indeterminate ('Nebsoy'; 'Improved Pelican')	
	, and the second of the second	
★ 18. MATURITY		
	GROUP: 000 2 = 00 3 = 0 4 = 1 5 = 11 6 = 111	λ = IΛ 8 = Λ
★ 18. MATURITY (	GROUP: 000 2 = 00 3 = 0 4 = 1 5 = 11 6 = 111 7 VI 10 = VII 11 = VIII 12 = IX 13 = X	z = IA 8 = A
★ 18. MATURITY (0 6 9 = 1)  ★ 19. DISEASE REA	GROUP:  000	' = IV 8 = V
★ 18. MATURITY (0 6 9 = 1)  ★ 19. DISEASE REA	GROUP: 000 2 = 00 3 = 0 4 = 1 5 = 11 6 = 111 7 VI 10 = VII 11 = VIII 12 = IX 13 = X	, = IA 8 = A
★ 18. MATURITY (  0 6 1 = 0 9 = 1	GROUP:  000	y = IV 8 = V
<ul> <li>★ 18. MATURITY (</li> <li>0 6 1 = 0</li> <li>9 = 1</li> <li>★ 19. DISEASE REALISE BACTERIAL</li> <li>★ 0 Bacterial</li> </ul>	GROUP:  000	y = IV 8 = V
<ul> <li>★ 18. MATURITY (1)</li> <li>★ 19. DISEASE REALISED BACTERIALISED BAC</li></ul>	GROUP:  000 2 = 00 3 = 0 4 = 1 5 = 11 6 = 111  VI 10 = VII 11 = VIII 12 = IX 13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)	v = IV 8 = V
★ 18. MATURITY (         0       6       1 = 0         9 = 1         ★ 19. DISEASE REAL         BACTERIAL         ★ 0       Bacte         ★ 0       Wildf	GROUP:  000 2 = 00 3 = 0 4 = 1 5 = 11 6 = 111  VI 10 = VII 11 = VIII 12 = IX 13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  (ire (Pseudomonas tabaci) ?	A = IA 8 = A
* 18. MATURITY (  0 6 1=0 9=1  * 19. DISEASE REAL  BACTERIAL  * 0 Bacte  * 0 Wildf  FUNGAL DISE	GROUP:  000  2 = 00  3 = 0  4 = 1  5 = 11  6 = 111  7  VI  10 = VII  11 = VIII  12 = IX  13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  fire (Pseudomonas tabaci) ?  EASES:	y = [V] 8 = V
★ 18. MATURITY 6         0       6       1 = 0         9 = 1         ★ 19. DISEASE REAL         BACTERIAL         ★ 0       Bacte         ★ 0       Wildf         FUNGAL DIST         ★ 0       Brown	GROUP:  000  2 = 00  3 = 0  4 = 1  5 = 11  6 = 111  7  VI  10 = VII  11 = VIII  12 = IX  13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  fire (Pseudomonas tabaci) ?  EASES:  In Spot (Septoria glycines)	y = [V] 8 = V
★ 18. MATURITY 6         0       6       1 = 0         9 = 1         ★ 19. DISEASE REAL         BACTERIAL         ★ 0       Bacte         ★ 0       Wildf         FUNGAL DIST         ★ 0       Brown	GROUP:  000  2 = 00  3 = 0  4 = 1  5 = II  6 = III  7  VI  10 = VII  11 = VIII  12 = IX  13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthornonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)	
* 18. MATURITY (  0 6 1 = 0 9 = 1  * 19. DISEASE REA  BACTERIAL  * 0 Bacte  * 0 Wildf  FUNGAL DISE  * 0 Brown  Froge  * 0 Race	GROUP:  000    2 = 00    3 = 0    4 = 1    5 = 11    6 = 111   7  VI    10 = VII    11 = VIII    12 = IX    13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  fire (Pseudomonas tabaci) ?  EASES:  In Spot (Septoria glycines)  eye Leaf Spot (Cercospora sojinal)	Other (Specify)
* 18. MATURITY (  0 6 1 = 0 9 = 1  * 19. DISEASE RE.  BACTERIAL  * 0 Bacte  * 0 Wildf  FUNGAL DISE  * 0 Brown  Froge  * 0 Race	GROUP:  000 2 = 00 3 = 0 4 = 1 5 = 11 6 = 111  VI 10 = VII 11 = VIII 12 = IX 13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  fire (Pseudomonas tabaci)  EASES:  In Spot (Septoria glycines)  eye Leaf Spot (Cercospora sojina)  1 Race 2 Race 3 Race 4 Race 5	
# 18. MATURITY (  0 6 1 = 0 9 = 1  # 19. DISEASE RE.  BACTERIAL  # 0 Bacte  # 0 Wildf  FUNGAL DISE  # 0 Brown  Froge  10 Target  10 Down	GROUP:  000  2 = 00  3 = 0  4 = 1  5 = II  6 = III  7  VI  10 = VII  11 = VIII  12 = IX  13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  fire (Pseudomonas tabaci) ?  EASES:  In Spot (Septoria glycines)  eye Leaf Spot (Cercospora sojina)  1  Race 2  Race 3  Race 4  Race 5	
★ 18. MATURITY         0       6       1 = 0         9 = 1         ★ 19. DISEASE READ         BACTERIAL         ★ 0       Bacte         ★ 0       Wildf         FUNGAL DISE         ★ 0       Brown         Froge         ★ 0       Race         0       Target         0       Powde	GROUP:  000  2 = 00  3 = 0  4 = 1  5 = II  6 = III  7  VI  10 = VII  11 = VIII  12 = IX  13 = X  ACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant)  L DISEASES:  erial Pustule (Xanthomonas phaseoli var. sojensis)  erial Blight (Pseudomonas glycinea)  iire (Pseudomonas tabaci)  2  EASES:  In Spot (Septoria glycines)  eye Leaf Spot (Cercospora sojinal)  1  Race 2  Race 3  Race 4  Race 5  t Spot (Corynespora cassiicola)  y Mildew (Peronospora trifoliorum var. manshurica)	

, 19. DISEASE REACT	ION: (Enter 0 = Not Tested; 1 = Suscept	ible; 2 = Resistant) (Continued)	
FUNGAL DISE.	ASES: (Continued)	rear Committee	9800001
★ 0 Pod and S	Stem Blight (Diaporthe phaseolorum var; s	sojae)	
	ed Stain <i>(Cercospora kikuchii)</i>		
0 Rhizoctor	nia Root Rot <i>(Rhizoctonia solani)</i>	/	
Phytophti	nora Rot (Phytophthora megasperma var.	sojae)	
★ 2 Race 1	O Race 2 2 Race 3	2 Race 4 2 Race	5 0 Race 6 0 Race 7
2 Race 8	0 Race 9 2 Other (Spe	, ,	Race 6 Race 7
VIRAL DISEASE	ES:		
0 Bud Blight	(Tobacco Ringspot Virus)	·.	
O Yellow Mo	saic (Bean Yellow Mosaic Virus)		
★ 0 Cowpea Mc	osaic (Cowpea Chlorotic Virus)		
0 Pod Mottle	(Bean Pod Mottle Virus)		
* 0 Seed Mottle	e (Soybean Mosaic Virus)		
NEMATODE DISI	EASES:		
Soybean Cy	rst Nematode (Heterodera glycines)	,	
★ 1 Race 1	0 Race 2 2 Race 3	0 Race 4 2 Other	(Specify) Race 14
0 Lance Nema	atode (Hoplolaimus Colombus)	on.a	Topachy)
★ 0 Southern Ro	oot Knot Nematode <i>(Meloidogyne incogn</i> i	ita)	
	oot Knot Nematode (Meloidogyne Hapla)	•	
O Peanut Root	Knot Nematode (Meloidogyne arenaria)		
0 Reniform Ne	ematode (Rotylenchulus reniformis)	·	
O OTHER DIS	EASE NOT ON FORM (Specify):		•
<b>→</b> [1] [	SPONSES: (Enter 0 = Not Tested; 1 = S	usceptible; 2 = Resistant)	
Iron Chlorosi	s on Calcareous Soil	,	
Other (Specif			
1 1	(Enter 0 = Not Tested; 1 = Susceptible; 2	? = Resistant)	
	Beetle (Epilachna varivestis)		
Totato Lear H	opper (Empoasca fabae)		
Other (Specify			
22. INDICATE WHICH VA	RIETY MOST CLOSELY RESEMBLES	THAT SUBMITTED.	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant Shape Leaf Shape	Essex	Seed Coat Luster	Mustang
Leaf Color	Mustang	Seed Size	Iroquois
Leaf Size	Mustang	Seed Shape	Resnik
:	Mustang	Seedling Pigmentation	Magellan
FORM LMGS-470-57 (6-83)			

VARIETY	NO. OF DAYS	PLANT LODGING	CM PLANT	LEAFL	ET SIZE	SEED COM	ITENT	SEED SIZE G/100	NO. SEEDS/
	MATURITY	SCORE	HEIGHT	CM Width	CM Length	% Protein	% Oil	SEEDS	POD
Submitted Mav <u>eri</u> ck	115	1.2	73.7	7.27	9.99	42	21	14	not recorde
I roquois Name of Similar Variety	113	1.6	88.9	7.09	9.75	41	21	14	not recorde

#### PUBLICATIONS USEFUL AS REFERENCE AIDS FOR COMPLETING THIS FORM:

- 1. Caldwell, B.E., ed. 1973. Soybeans: Improvement, Production, and Uses. Amer. Soc. Agron. Monograph No. 16.
- 2. Buttery, B.R. and R.I. Buzzell. 1968. Peroxidase activity in seeds of soybean varieties. Crop Sci., 8: 722-725.
- 3. Hymowitz, T. 1973. Electrophoretic analysis of SBTI-A2 in the USDA soybean germplasm collection. Crop Sci., 13: 420-421.
- 4. Payne, R.C. and L.F. Morris. 1976. Differentiation of soybean cultivars by seedling pigmentation patterns. J. Seed Technol. 1: 1-19.

#### Exhibit D

#### Additional Description of the Variety

At soybean cyst nematode infested locations, seed yield of Maverick was 25% higher than Iroquois and 3% higher than Jack. Compared with Iroquois at noninfested soybean cyst nematode sites, Maverick was similar in yield. Maverick is similar in seed quality score and seed oil concentration as Iroquois. This data was compiled from USDA Northern Regional Tests.

### SCN UNIFORM TEST III 1996

			Previous
Strain			testing
1 Jack	Fayette	x Hardin	11
2 Iroquois (III)	LŇ81-1029	x Asgrow A2943	. 1
3 IA3005 `	L82C-1212	x Kenwood	3
4 A94-773014	Pioneer P9303	x A87-395012	95 SCN PIII
5 A94-773038	Jack	x Pioneer P9303	95 SCN PIII
6 LN92-10484	Jack	x A86-303014	95 SCN PIII
7 LN92-10855	Jack	x Asgrow A3205	95 SCN PIII
8 Maverick	LN86-4668	x Resnik	1
IA3005 =	A92-726034		
Maverick =	SS95-1000 reselecti	on of SL90-4455	

SCN UNIFORM TEST III 1996

	<u>. 1</u>	2CIN DIV	ITURW I		1990	 	
	Tipp	. Co.	Pulas	ki Co.			
	Ī		$\mathbf{n}$	1			
Strain	Ag. Al	lumni	Sto	otler		 	
	%	React	%	React			
1 Jack	26.9	S	1.4	R			
2 Iroquois (III)	100.0	S	100.0	S			
3 IA3005	13.0	S	9.0	R			
4 A94-773014	38.5	S	9.7	R			
5 A94-773038	11.1	S	5.6	R			
6 LN92-10484	56.7	S	4.2	R			
7 LN92-10855	13.0	S	13.9	. <b>S</b>			
8 Maverick	40.4	S	26.4	S			

Indiana greenhouse tests, refer to page vi of Methods.

SCN UNIFORM TEST III 1996

Strain	Iron Chlorosis Humboldt	Urb	nois ana ce 3		SDS Ridgway IL	
	IA	vigor	root	incidence	severity	index
	score	score	score	%	score	
l Jack	4.0	3.5	1.0	3.2	1.5	0.48
2 Iroquois (III)	4.4	3.5	2.5	53.9	1.2	6.84
3 IA3005	3.2	1.5	0.5	0.0	1.0	0.00
4 A94-773014	4.2	1.0	0.5	40.3	1.3	4.61
5 A94-773038	3.9	2.5	0.0	23.7	1.1	2.56
6 LN92-10484	3.8	1.5	0.0	36.2	1.6	6.42
7 LN92-10464	4.6	1.5	0.0	8.8	1.0	0.69
8 Maverick	4.0	1.5	0.0	6.3	1.2	0.78

NI non-infested; I infested

Iowa Iron Chlorosis rating, refer to page vi Methods.

Illinios SCN vigor and root ratings, refer to page vi Methods.

Illinois SDS evaluation, refer to page vi Methods

Illinois Sudden Death Syndrome rating: Plots are scored at plant stage R6, Ridgway, IL, Southern Illinois University. All disease scores were interpolated to the R=6.2 reproductive stage. Some scores are <0 or >100 due to the Lattice adj.

R6DI = SDS Disease Incidence (% of plants with visible symptoms).

R6DS = SDS Disease Severity (1 = mild chlorosis, 5 = severe leaf scorch, 9 = premature plant death).

R6DX = SDS Disease Index (R6DI x R6DS/9)

*Indiana's greenhouse test:* Soil from each field location is brought to the laboratory and used to test each SCN soybean line for resistance to the SCN population found in that field. Seed of each soybean line are germinated in sand. When seedlings are several inches tall, sand is washed from the roots. Each seedling is placed in a 1-pint plastic cup partially filled with a soil:sand (1:2) mixture, 1 ml of inoculum is pipetted over the roots and additional soil/sand mixture is added to the pot. Five replicates of each entry are set up in this fashion. Inoculum is prepared by extracting cysts from the field soil, dissolving the cyst cuticle with sodium hypochlorite to release eggs and juveniles, and adjusting the inoculum concentration to 2000-3000 per ml. Plants are grown for a period of 6-8 weeks at a temperature around 75 F. When it is judged that development of first generation SCN females has taken place, tops are cut off the plants, and the roots and soil washed to recover all females. Total numbers recovered from each replicate of each entry are used to calculate an index of parasitism, with the mean number of cysts developing on the susceptible entries equal to 100%. If the mean number of cysts on a regional entry is less than 10% of the number on the susceptible varieties, the entry is rated as resistant. In addition to a S or R notation, the actual percentage is reported.

Maryland SCN rating: Number of cysts = full cysts/4 plants sampled on July 17, 1996.

Illinois SCN rating: SCN root score based on 0 = 0 cysts; 1 = 1-5 cysts; 2 = 6-10 cysts; 3 = 11-30 cysts; and 4 > 30 cysts per plant (average of 5 plants per replication). SCN vigor score based on 1=no stunting or plant death to 5=all plants dead.

<u>Iron chlorosis</u> is rated from 1= no chlorosis, to 5= severe chlorosis.

system (Golden et al., 1970; Schmitt and Shannon, 1992), and was calculated as follows: reflect the number of females and cysts per plant for three replications. Data were collected at the University of Missouri-Columbia. The Reaction of Maverick to Race 3 and Race 14 of the soybean cyst nematode in comparison to the susceptible variety Lee. The data index of parasitism (IP%) was used as the criteria to distinguish resistance based on the standard soybean cyst nematode classification

number of females and cysts on a given individual

P(%) = ----- x 100

average number of females and cysts on Lee

			Race 3				Race 1	14		
Variety	Numl	ber of c	Number of cysts per plant	Mean	Mean Index of	Number	er of c	of cysts per plant	Mean	Mean Index of
	Rep	l Rep	Rep 1 Rep 2 Rep 3		Parasitism	Rep 1 R	Rep 2	tep 2 Rep		<b>Paratism</b>
Maverick	5	0	4	3	4 %	24	30	13	22	20%
Lee	75	70	60	68		101	118	108	109	
								7-500-		

# References

of the soybean cyst nematode (Heterodera glycines). Plant Dis. Rep. 54:544-546. Golden, A.M., J.M. Epps, R.D. Riggs, L.A. Duclos, J.A. Fox, and R.L. Bernard. 1970. Terminology and identity of infraspecific forms

Schmitt, D.P. and G. Shannon. 1992. Differentiating soybean responses to Heterodera glycines Races. Crop Sci. 32:275-277.

2	CALLY. Include form number and date on all reproductions.  U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	l ine iouowing statements are made	FORM APPROVED - OMB NO. 0581-00
		To the dialor desay one trap	erwork Reduction Act (PRA) of 1995 .
ST	EXHIBIT E ATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to certificate is to be issued (7 U.S.C. until certificate is issued (7 U.S.C.	o determine if a plant variety protection 2421). Information is held confidenti 2426).
1. NAME OF APPL		2. TEMPORARY DESIGNATION	3. VARIETY NAME
The Curat	cors of the University of Missouri	OR EXPERIMENTAL NUMBER	
-		LN90-4455	Maverick
4. ADDRESS (Stre	eet and No., or R.F.D. No., City, State, and ZIP Code, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
Universit	y of Missouri	573-882-3211	573-884=5446
311 Jesse		7. PVPO NUMBER	373-064-3440
Columbia,	MO 02511		a
8. Does the appl	licant own all rights to the variety? Mark an "X" in appropriat	e block. If no please explain	*
	у предпас	e block. If no, please explain.	X YES NO
\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		,	
9. Is the applican	t findividual or company) a U.S. national or U.S. based, compa	ny?	
If no, give nan	ne of country		X YES NO
	YES NO If no, give name of country  f original rights to variety were owned by a company, is the o  YES NO If no, give name of country  Dianation on ownership (If needed, use reverse for extra space)	riginal owner(s) a U.S. based compa	· <del></del>
PLEASE NOTE:			
	tion can be afforded only to owners (not licensees) who meet		
of a country whi	ne variety are owned by the original breeder, that person mus ch affords similar protection to nationals of the U.S. for the s	it be a U.S. national, national of a U ame genus and species.	POV member country, or national
<ol> <li>If the rights to the nationals of a UF genus and specie</li> </ol>	ne variety are owned by the company which employed the ori POV member country, or owned by nationals of a country whees.	ginal breeder(s), the company must ich affords similar protection to nat	be U.S. based, owned by fonals of the U.S. for the same
. If the applicant is	s an owner who is not the original owner, both the original ow	vner and the applicant must meet or	ne of the above criteria.
	owner may be the individual or company who directed final b	· ·	
ollection is estimate	perwork Reduction Act of 1995, no persons are required to be valid OMB control number for this information collection and to average 10 minutes per response, including the time f data needed, and completing and reviewing the collection of t	is 0581-0055. The time requi	
he U.S. Departmen Isability, political be Iternative means for 1 (202) 720-5881 (v	of of Agriculture (USDA) prohibits discrimination in its progreeliefs, and marital or familial status. (Not all prohibited by a communication of program information (braille, large print, a communication) 720-7808 (TDD).  Write the Secretary of Agriculture, U.S. Department of Agricu	ams on the basis of race, color, rases apply to all programs.) Personautionary etc.) should contact the	USDA Office of Communications